

FEDERAL PUBLIC SERVICE COMMISSION



COMPETITIVE EXAMINATION FOR RECRUITMENT TO POSTS IN BS-17 UNDER THE FEDERAL GOVERNMENT, 2012

Roll Number

PURE MATHEMATICS, PAPER-I

TIME ALLOWED: THREE HOURS

MAXIMUM MARKS: 100

- NOTE:**
- (i) Candidate must write **Q. No.** in the **Answer Book** in accordance with **Q. No.** in the **Q. Paper.**
 - (ii) Attempt **FIVE** questions in all by selecting **THREE** questions from **SECTION-A** and **TWO** questions from **SECTION-B.** **ALL** questions carry **EQUAL** marks.
 - (iii) Extra attempt of any question or any part of the attempted question will not be considered.
 - (iv) **Use of Scientific Calculator is allowed.**

SECTION-A

- Q. 1.** (a) Let H be a normal subgroup and K a subgroup of a group G . Prove that HK is a subgroup of G and $H \cap K$ is normal in K and $\frac{HK}{H} \cong \frac{K}{H \cap K}$. (12)
- (b) Show that number of elements in a Conjugacy class Ca of an element 'a' in a group G is equal to the index of its normaliser. (8)
- Q. 2.** (a) Prove that if G is an Abelian group, then for all $a, b \in G$ and integers n , $(ab)^n = a^n b^n$. (6)
- (b) Show that subgroup of Index 2 in a group G is normal. (7)
- (c) If H is a subgroup of a group G , let $N(H) = \{a \in G \mid aHa^{-1} = H\}$ Prove that $N(H)$ is a subgroup of G and contains H . (7)
- Q. 3.** (a) Show that set C of complex numbers is a field. (6)
- (b) Prove that a finite integral domain is a field. (6)
- (c) Show that $\bar{Z}_6 = \{\bar{0}, \bar{1}, \bar{2}, \bar{3}, \bar{4}, \bar{5}\}$ is a ring under addition mod 6 and multiplication mod 6 but not a field. Find the divisors of Zero in \bar{Z}_6 . (8)
- Q. 4.** (a) Let F be a field of real numbers, show that the set V of real valued continuous functions on the closed interval $[0,1]$ is a vector space over F and the subset Y of V containing all functions whose n th derivatives exist, forms a subspace of V . (10)
- (b) Prove that any finite dimensional vector space is isomorphic to F^n . (10)
- Q. 5.** (a) State and prove Cayley-Hamilton theorem. (10)
- (b) Use Cramer's rule to solve the following system of linear equations: (10)
- $$\begin{aligned} x + y + z + w &= 1 \\ x + 2y + 3z + 4w &= 0 \\ x + y + 4z + 5w &= 1 \\ x + y + 5z + 6w &= 0 \end{aligned}$$

SECTION-B

- Q. 6.** (a) Prove that an equation of normal to the astroid $x^{2/3} + y^{2/3} = a^{2/3}$ can be written in the form: (10)
- $$y \cos \theta - x \sin \theta = a \cos 2\theta$$
- Hence show that the evolute of the curve is
- $$(x + y)^{2/3} + (x - y)^{2/3} = 2a^{2/3}$$
- (b) If r_1 and r_2 are radii of curvature at the extremities of any chord of the Cardioid $r = a(1 + \cos \theta)$ which passes through the pole, then prove that (10)
- $$r_1 r_2 = \frac{16a^2}{9}$$

PURE MATHEMATICS, PAPER-I

- Q. 7. (a)** Find an equation of the normal at any point of the curve with parametric equations: **(10)**
 $x = a(\cos t + t \sin t)$, $y = a(\sin t - t \cos t)$.
Hence deduce that an equation of its evolute is $x^2 + y^2 = a^2$.

- (b)** Find equations of the planes bisecting the angle between the planes **(10)**
 $3x + 2y - 6z + 1 = 0$ and $2x + y + 2z - 5 = 0$.

- Q. 8. (a)** Define a surface of revolution. Write equation of a right elliptic-cone with vertex at **(6)**
origin.

- (b)** Identify and sketch the surface defined by **(6)**
 $x^2 + y^2 = 2z - z^2$.

- (c)** If $y=f(x)$ has continuous derivative on $[a, b]$ and S denotes the length of the arc of **(8)**
 $y=f(x)$ between the lines $x=a$ and $x=b$, prove that

$$S = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} .$$

Find the length of the parabolas $y^2 = 4ax$

- (i) From vertex to an extremity of the latus rectum.
(ii) Cut off by the latus rectum.

FEDERAL PUBLIC SERVICE COMMISSION



COMPETITIVE EXAMINATION FOR RECRUITMENT TO POSTS IN BS-17 UNDER THE FEDERAL GOVERNMENT, 2012

Roll Number

PURE MATHEMATICS, PAPER-II

TIME ALLOWED: THREE HOURS

MAXIMUM MARKS: 100

- NOTE:**
- (i) Candidate must write **Q. No.** in the **Answer Book** in accordance with **Q. No.** in the **Q. Paper.**
 - (ii) Attempt **FIVE** questions in all by selecting **THREE** questions from **SECTION-A** and **TWO** questions from **SECTION-B.** **ALL** questions carry **EQUAL** marks.
 - (iii) Extra attempt of any question or any part of the attempted question will not be considered.
 - (iv) **Use of Scientific Calculator is allowed.**

SECTION-A

- Q. 1. (a)** State and prove Taylor's theorem with Cauchy's form of remainder. (8)
- (b)** Evaluate (i) $\lim_{x \rightarrow 0} \left(\frac{1}{x}\right)^{\tan x}$ (ii) $\int e^{ax} \sin(bx + c) dx$ (6)
- (c)** Show that $\int_0^{\pi/2} \sin^p x \cos^q x dx = \frac{\Gamma\left(\frac{p+1}{2}\right)\Gamma\left(\frac{q+1}{2}\right)}{2\Gamma\left(\frac{p+q}{2} + 1\right)}$ (6)
- Q. 2. (a)** Sketch the graph of the curve $r^2 = a \sin 2\theta, a > 0$. Also write pedal equation for this curve. (8)
- (b)** Show that the parabola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ has asymptotes $y = \frac{b}{a}x$ and $y = -\frac{b}{a}x$ (6)
- (c)** Define extrema (local and global) of a function of two variables. Find three positive numbers whose sum is 48 and whose product is as large as possible. (6)
- Q. 3. (a)** Find the volume of the tetrahedron bounded by the coordinate planes and the plane $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1, a, b, c > 0$. (8)
- (b)** Evaluate $\int_0^{\pi/2} \ln(\sin x) dx$ (6)
- (c)** Determine the values of x for which the power series $\sum_{n=2}^{\infty} \frac{x^n}{\ln n}$ converges absolutely, converges conditionally and diverges. (6)
- Q. 4. (a)** Define a metric on a non-empty set X . If d is a metric on X , show that if $d'(x, y) = \frac{d(x, y)}{1 + d(x, y)}$ then d' is also a metric on X . Also write open and closed balls (spheres) in the discrete metric space (X, d) with radius 1 and 1.1 centered at some $x \in X$. (5+3+2=10)
- (b)** Define limit point of a subset A of a metric space X . Show that an open sphere containing a limit point x of A contains infinitely many points of A other than x . (10)

PURE MATHEMATICS, PAPER-II

- Q. 5. (a)** Show that R^n is a complete metric space under the metric defined by (7)
 $d(x, y) = \sqrt{\sum (\xi_i - \eta_i)^2}$, $x, y \in R^n$
 Where $x = (\xi_1, \xi_2, \dots, \xi_n)$ and $y = (\eta_1, \eta_2, \dots, \eta_n)$
- (b)** Show that a function $f: (X, d) \rightarrow (Y, d')$ is continuous if and only if for an open subset V (7)
 of Y, $f^{-1}(V)$ is an open subset of X.
- (c)** Find the radius of convergence and interval of convergence of the power series: (6)

$$\sum_{n=0}^{\infty} \frac{(-1)^{n+1} (x+1)^{2n}}{(n+1)^2 5^n}$$

SECTION-B

- Q. 6. (a)** If C is a continuous curve and $f(z)$ is defined on each point of C, then prove that (10)

$$\left| \int_C f(z) dz \right| \leq ML$$

 Where $M = \max |f(z)|$ and L is length of curve C.
- (b)** Suppose $f(z) = U(x, y) + iV(x, y)$ is differentiable at a point $z = x + iy$, then at z the (10)
 first order partial derivatives of U and V exist and satisfy Cauchy-Reiman equations:

$$\frac{\partial U}{\partial x} = \frac{\partial V}{\partial y}, \frac{\partial U}{\partial y} = -\frac{\partial V}{\partial x}.$$

 Verify Cauchy-Reiman equations for the function $f(z) = e^{-x} \cos y - i e^{-x} \sin y$.
- Q. 7. (a)** Define singularity of a function $f(z)$. Investigate for the pole, singularities and zeros, (6)
 the function $f(z) = z^2$
- (b)** Let D be simply connected domain and $f(z)$ be analytic in D. Let $f'(z)$ exist and is (6)
 continuous at each point of D then prove that $\int_C f(z) dz = 0$, where C is any closed
 Contour in D.
- (c)** State De Moivre's theorem and hence prove that (8)
 (i) $\cos 5\theta = 16 \cos^3 \theta - 20 \cos^2 \theta + 5 \cos \theta$
 (ii) $\sin^n \theta = (-1)^{\frac{n-1}{2}} \frac{1}{2^{n-1}} \left[\sin n\theta - \sin(n-2)\theta + \frac{n(n-1)}{2} \sin(n-4)\theta - \dots \right]$
- Q. 8. (a)** Solve the equation $x^{12} - 1 = 0$ and find which of its roots satisfy the equation $x^4 + x^2 + 1 = 0$. (6)
- (b)** Show that multiplication of a vector z by $e^{i\alpha}$ where α is a real number, rotates the (6)
 vector z counter clockwise through an angle of measure α .
- (c)** Sum the series (8)

$$n \sin \theta + \frac{n(n+1)}{2!} \sin 2\theta + \frac{n(n+1)(n+2)}{3!} \sin 3\theta + \dots$$
