



**FEDERAL PUBLIC SERVICE COMMISSION
COMPETITIVE EXAMINATION FOR
RECRUITMENT TO POSTS IN BPS-17 UNDER
THE FEDERAL GOVERNMENT, 2009**

S.No.	
R.No.	

STATISTICS

TIME ALLOWED:	(PART-I) 30 MINUTES	MAXIMUM MARKS:20
	(PART-II) 2 HOURS & 30 MINUTES	MAXIMUM MARKS:80

- NOTE:** (i) First attempt **PART-I (MCQ)** on separate **Answer Sheet** which shall be taken back after **30 minutes**.
(ii) **Overwriting/cutting of the options/answers will not be given credit.**
(iii) **Statistical Table will be provided if required.**
(iv) **Use of Scientific Calculator is allowed.**

PART – I (MCQs)

(COMPULSORY)

Q.1. Select the best option/answer and fill in the appropriate box on the Answer Sheet. (20)

- (i) The probability of event given the event B is $P(A/B)$ is equal to $P(A)$ if B is:
(a) any event in sample S (b) sample space S
(c) $A \subset B$ (d) B is dependent on A
- (ii) If an event $A = (A \cap B_1) \cup (A \cap B_2) \cup \dots \cup (A \cap B_n)$ and sample space $S = B_1 \cup B_2 \cup \dots \cup B_n$ and $B_i \cap B_j = \emptyset$, $i \neq j$, $i, j = 1, 2, \dots, n$ then:
(a) $P(A)=1$ (b) $P(A) = \sum_{i=1}^n P(B_i)$
(c) $P(A) = \sum_{r=1}^n P(A|B_r)P(B_r)$ (d) $P(A) = \sum_{i=1}^n P(B_i|A)$
- (iii) A family has two children, then the probability of the event that atleast one of them is a boy is:
(a) $\frac{1}{4}$ (b) $\frac{1}{2}$
(c) $\frac{1}{3}$ (d) $\frac{3}{4}$
- (iv) The value of $\sum_{k=0}^n \binom{n}{k}$ is:
(a) n^k (b) k^n (c) 2^k (d) 2^n
- (v) A student is attempting to log on internet with 0.5 chance of successful attempt in each trial. The average number of attempts required to log on successfully is:
(a) 1 (b) 2 (c) 3 (d) 4
- (vi) The mean of binomial random variable, with parameter probability of success is twice the probability of failure in a single trail then:
(a) greater than $\frac{2}{9} \cdot n$ (b) less than twice the variance (c) greater than $\frac{2n^2}{3}$ (d) none of these
- (vii) If x follows normal distribution with pdf $\exp - \bar{\lambda} x^2$ then its mean and variance are:
(a) Π, Π (b) $\bar{\lambda}, \bar{\lambda}/2$ (c) $\sigma, \frac{1}{2\bar{\lambda}}$ (d) 0, 1
- (viii) Let X be the number of patients arriving at OPD on any day in a hospital according to poisson distribution with the probability of at least one arrival in a day is $1 - e^{-2}$. Then average number of arrivals of patients per day is:
(a) 8 (b) 4 (c) 2 (d) 1
- (ix) To test the hypothesis $H_0: \mu_1 = \mu_2 = \mu_3$ at $\alpha = 0.05$, then one can use:
(a) Regression Analysis (b) Analysis of Variance (c) z-test (d) t-test

STATISTICS

- (x) Suppose we have random sample of size n from normal population with mean μ and variance σ^2 , then maximum likelihood estimate of σ^2 , when $\hat{\mu} = \bar{x}$, is:
- (a) $\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$ (b) $\frac{1}{n} \sum_{i=1}^n (x_i - \mu)^2$ (c) $\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$ (d) $\frac{1}{n} \sum_{i=1}^n (x_i - \mu)^2$
- (xi) The probability of accepting a hypothesis when it is false is 0.2 then the probability of rejecting this hypothesis when it is false is:
- (a) 0.95 (b) 0.9 (c) 0.85 (d) 0.8
- (xii) If $(x_1, y_1), \dots, (x_n, y_n)$ set of n observations on Variable $X =$ Hours studied, random variable $Y =$ test score and $Y = a + bx$ is the least square line that approximates the regression of test scores on the number of hours studied is given by $Y = 21.819 + 3.471 X$. If the desired test score is at least 60 then hours of studied should be at least:
- (a) none (b) at most 10 (c) at least 10 (d) at least 11
- (xiii) The inter arrival time between two messages in a communication/service system follows negative exponential distribution $2e^{-2x}, x > 0$, then average inter arrival time between two messages is:
- (a) 1 (b) 2 (c) $\frac{1}{2}$ (d) $\frac{1}{4}$
- (xiv) In random sampling with replacement, the probability that all n specified units of a sampling frame are selected in n draws, with population size is N , is:
- (a) $\frac{1}{n^N}$ (b) $\frac{1}{N^n}$ (c) $\frac{1}{n}$ (d) $\left(\frac{1}{N}\right)^n$
- (xv) For population with heterogeneous groups, the suitable sampling scheme is:
- (a) Simple Random Sampling (b) Systematic Sampling
(c) Cluster Sampling (d) Stratified Sampling
- (xvi) The variance of x, y, z, u, v objects is:
- (a) 5 (b) $\sqrt{5}$ (c) 1 (d) none of these
- (xvii) For a size of size n from $N(\mu, \sigma^2)$, σ^2 is unknown, $H_0: \mu = \mu_0$ against $H_1: \mu \neq \mu_0$ then:
- (a) t test with $n - 1$ d.f. at $\alpha = 0.05$ will be used
(b) F test with n d.f. at $\alpha = 0.05$ will be used
(c) t test with $n - 1$ d.f. at $\alpha = 0.025$ will be used
(d) X^2 test with $n - 1$ d.f. at $\alpha = 0.025$ will be used
- (xviii) Height of date trees, say, follow $N(8, 4)$ then the third moment about mean is:
- (a) 3×64 (b) 4×256 (c) 0×4 (d) none of these
- (xix) In a sample of size n , x are girls with variance $V(x)$ and $n-x$ are boys with variance is:
- (a) $V(x)$ (b) $V(x) + n^2$ (c) $V(x) - n^2$ (d) none of these
- (xx) If two random variables are independent then correlation or covariance zero. If correlation or covariance between two variables X and Y is zero then:
- (a) X and Y are independent of one another (b) X and Y may be independent on one another
(c) X and Y may be mutually exclusive (d) None of these

PART – II

NOTE:	<p>(i) PART-II is to be attempted on the separate Answer Book.</p> <p>(ii) Attempt ONLY FOUR questions from PART-II. All questions carry EQUAL marks.</p> <p>(iii) Extra attempt of any question or any part of the attempted question will not be considered.</p>
--------------	---

- Q.2.** (a) Explain the concept of conditional probabilities using daily life events. Also justify the common formula of conditional probability of an event A given B is $P(A|B) = \frac{P(A \text{ and } B)}{P(B)}$. (4)
- (b) In answering a question on a MCQ test a student either knows the answer or guesses. Let p be the probability that student knows the answer and $1-p$ the probability that he/she guesses. Assume that a student who guesses at the answer will be correct with probability $\frac{1}{n}$, where n is the number of MC alternatives. What is the conditional probability that a student knew the answer to a question given that she answered it correctly? (8)

STATISTICS

- (c) A Laboratory blood test is 95% effective in detecting a certain disease when it is, in fact, present but also yields a “false positive” result for 1% of the healthy persons tested. If 0.5 percent of the population actually has the disease, what is the probability a person has the disease given that his test result is positive. (8)
- Q.3.** If X is the amount of money (in Hundreds of rupees) that a salesman spends on gasoline during a day and Y is the corresponding amount of money (in Hundred of rupees) for which he/she is reimbursed, the joint density of these two random variables is given by $f(x, y) = K \cdot \frac{20-x}{x}$, for $10 < x < 20$, $\frac{x}{2} < y < x$ and 0 elsewhere. Find
- (a) K (b) $f_x(x)$ (c) $f_{Y|X}(y|x=12)$
- (d) the probability that the salesman will be reimbursed at least 8 units of money when spending 12 units of money. (4×5)
- Q.4.** In a certain city three T.V. channels are available. During prime time on Saturday nights Channel 1 has 50% of the viewing audience, Channel 2 has 25% of the viewing audience and Channel 3 has remaining percent of the viewing audience:
- (a) Compute the probability that among 10 T.V. viewers in that city, randomly chosen on a Saturday night, 50% watching Channel 1, 30% watching Channel 2 and 20% watching Channel 3. (10)
- (b) Calculate the average number of viewers watching Channel 1, Channel 2, Channel 3 out of 10 randomly selected. (10)
- Q.5.** The best yardstick to measure the social and moral maturity of a society is the state of its children. In a recent report titled ‘The State of Pakistan Children 2007’ the infant mortality rate at 84 per 1000 live births, under-five mortality rate is 125 per 1000 and 38% of children under five being under-weight.
- (a) Construct 95% C.I. for infant mortality rate. (5)
- (b) Construct 95% C.I. for under-five mortality rate. (5)
- (c) Construct 95% C.I. for children under-five being under weight. (5)
- (d) Write a brief report in the light of inferences made in (a), (b) and (c) so that non-technical person can understand. (5)
- Q.6.** (a) Define Chi-square Goodness-of-fit test with a simple example. (8)
- (b) Mendelian theory indicates that the shape and colour of certain variety of pea ought to be grouped into 4 groups, “round and yellow,” “round and green,” “angular and yellow” and “angular and green,” according to ratio 9/3/3/1. For a sample of size $n = 556$ peas, the following results were obtained: Round and Yellow 315, Round and green 108, Angular and yellow 101 and Angular and green 32. Test $H_0: p_1 = \frac{9}{16}, p_2 = \frac{1}{3}, p_3 = p_2$ and $p_4 = 1 - p_1 - p_2 - p_3$. (12)
- Q.7.** (a) To learn good programming techniques, two courses: C++ and C-Sharp are taught by an I.T Department of a University. The success of each course is evaluated by the scores achieved by the students in the Departments Programmers Test. Nine students using course C++ achieved an average test score of 89.6 with a sample variance of 12.96. Seven students using course C-Sharp got an average score of 81.9 with a sample variance of 161.29. Assuming all test scores are normally distributed, test $H_0: \mu_x = \mu_y$ against $H_a: \mu_x > \mu_y$ at $\alpha = 0.01$. (8)
- (b) (i) Explain a test statistic which test the hypothesis on difference of two variances on normal populations. (6)
- (ii) Consider part (a) of Q.7. At $\alpha = 0.05$, whether it is reasonable to assume that the variance is the same for two courses mentioned above. (6)
- Q.8.** (a) Explain Systematic Sampling with an example. Compare this method of sampling with simple random sampling. (6)
- (b) Describe the relationship of systematic sampling with Cluster Sampling (6)
- (c) Write notes on the following terms: (8)
- (i) Maximum Likelihood Estimation (ii) Least Squares Estimation of Regression Coefficient.
- (iii) Census and Registration (iv) Bayes Theorem
